E4-2025-4

A. D. Efimov^{1,2,*}, I. V. Koval¹, I. N. Izosimov^{**}

COMPARISON OF THE DESCRIPTION OF YRAST BANDS IN THE HARRIS AND **IBM** MODELS FOR EVEN NUCLEI TH AND U

Submitted to "Chinese Physics C"

¹ Admiral Makarov State University of Maritime and Inland Shipping, Saint Petersburg, Russia

² Ioffe Physical-Technical Institute, Russian Academy of Sciences, Saint Petersburg, Russia

^{*} E-mail: efimov98@mail.ru

^{**} E-mail: izosimov@jinr.ru

Ефимов А. Д., Коваль И. В., Изосимов И. Н. Сравнение описания ираст-полос в моделях Харриса и IBM для четных ядер Th и U

Для четных ядер Th и U изучено описание энергий ираст-полос в модели переменного момента инерции Харриса в ее феноменологическом применении. Для некоторых из этих ядер приводятся результаты, полученные в микроскопической версии IBM, в которой рассматриваются и высокоспиновые моды возбуждений. Для других ядер в тех случаях, когда высокоспиновые орбиты не столь существенны при описании свойств ираст-полос, помимо модели Харриса применяется IBM с полным стандартным гамильтонианом, построенным только из s-, d-бозонов. Так как модель Харриса рассматривается не более чем с четырьмя параметрами, она как для тяжелых, так и сверхтяжелых ядер может быть использована для аппроксимации энергий высокоспиновых состояний. Исследование свойств ядер в различных моделях позволяет выявить возможности описания различного поведения моментов инерции в IBM и наметить дополнительные критерии относительно реализации пересечения полос.

Работа выполнена в Лаборатории ядерных реакций им. Г. Н. Флерова ОИЯИ.

Препринт Объединенного института ядерных исследований. Дубна, 2025

Efimov A. D., Koval I. V., Izosimov I. N.

E4-2025-4

E4-2025-4

Comparison of the Description of Yrast Bands in the Harris and IBM Models for Even Nuclei Th and U

For even Th and U nuclei, the description of the energies of yrast bands in the Harris variable moment of inertia model is studied in its phenomenological application. For some of these nuclei, the results obtained in the IBM microscopic version are presented, in which high-spin excitation modes are also considered. For other nuclei, in cases where high-spin orbits are not so significant in describing the properties of yrast bands, in addition to the Harris model, IBM with a full standard Hamiltonian constructed only from *s*, *d* bosons is utilized. Since the Harris model is considered with no more than four parameters, it can be used for both heavy and superheavy nuclei to approximate the energies of high-spin states. The study of the properties of nuclei in various models will reveal the possibilities of describing different behavior of the moments of inertia in IBM and outline additional criteria for the implementation of band crossing.

The investigation has been performed at the Flerov Laboratory of Nuclear Reactions, $\ensuremath{\mathsf{JINR}}$.

INTRODUCTION

Visualization of the behavior of the energies of states E in a band becomes more qualitative if we move from energies to effective moments of inertia J. This also allows us to improve the method of comparing experimental and calculated energies of states of one band.

In addition, the backbending effect can be used to form judgments about the spin values at which the bands cross. Comparison of the $J(\omega^2)$ curves obtained from experimental energies with those given by various nuclear models, including microscopic ones, can reveal the capabilities of the corresponding models and provide more reasoned judgments about the nature and character of the states.

One of the methods of reproducing and predicting energies is based on the expansion of the moment of inertia in powers of the rotation frequency ω , as in the Harris model [1]. This is justified when the transition to a new band has not yet occurred in the rotation band. As a rule, such a transition occurs quite quickly in one or two states of the yrast band. In this case, this is shown on the graph of the moment of inertia versus the square of the rotation frequency in a specific way through backbending. Among heavy nuclei, starting with thorium isotopes, three such nuclei are known today: 220 Th, 242 Pu, and 244 Pu. This is why in [2], within the framework of the IBM1 [3] phenomenology (hereinafter referred to as IBM), a good description of the energies of yrast bands up to extremely high spins *I* in heavy nuclei from Pu to No was obtained. The effective moment of inertia and the square of the rotational speed are defined as

$$\frac{2J}{\hbar^2} = \frac{4I-2}{E(I \to I-2)}, \quad (\hbar\omega)^2 = \frac{\left(E(I \to I-2)\right)^2}{\left(\left(I(I+1)\right)^{1/2} - \left((I-2)(I-1)\right)^{1/2}\right)^2}.$$
 (1)

This paper presents a description of the energies of yrast bands in even-even heavy nuclei in the Harris model, considered not from the point of view of the cracking model, but as a phenomenological model with a number of parameters no more than four. The possibility of IBM in its phenomenological aspect to describe moments of inertia is also considered. The moments of inertia obtained in the two models are compared with each other. Finally, the task is set to develop a criterion for determining the spin at which the crossing of bands occurs. This can be done based on a comparison of the behavior of $J(\omega^2)$ according to experimental data and what the Harris model gives. It should be noted that in some cases, although backbending may not

be observed, the crossing of bands is still possible. This was revealed by microscopic description of the chain of isotopes $^{220-236}$ Th [4].

In order to increase the calculated moment of inertia at high-spin states in deformed nuclei (162,164 Er), in [5], a new term was incorporated into the Hamiltonian of IBM1, so that the operator $\hat{I}^2 \rightarrow \hat{I}^2/(1+f\hat{I}^2)$. This part of the Hamiltonian is always diagonal for any values of the parameters $H_{\rm IBM}$. This addition is motivated by the fact that a decrease in superfluidity, the gap parameter, leads to an increase in the moment of inertia, and the gap parameter decreases with increasing rotation frequency. This should lead to dJ/dI > 0 and $dJ/d\omega > 0$. Here, however, it should be kept in mind that the growth of the moment of inertia from the rotation frequency is achieved with arbitrary values of the traditional and full Hamiltonian of IBM, except for the one that corresponds to the case of the SU(3) limit of IBM. In addition, with the growth of the spin of the collective state, the influence of two-quasiparticle states on the energy of this state increases. One example of a band built on a two-quasiparticle mode is the S band. This also leads to an additional increase in the moment of inertia or a decrease in the energies of intra-band transitions.

Analysis of the dynamic moments of inertia of the lowest superdeformed (SD) bands in even-even Hg, Pb, Gd, and Dy [6] and 192,194 Hg, 152,154 Dy [7] isotopes led to the need to extend the coefficient *f* so that considering only the relative excitation of the states in a rotational band, the energy of the state with angular momentum *I* can be simply expressed as

$$E(I) = \frac{C_0}{1 + f_1 I(I+1) + f_2 I^2 (I+1)^2} I(I+1).$$

Introduction of an additional modification of the Arima coefficient f shows that J can change nonmonotonously with rotational frequency. Up to the point that as the spin increases, the positive slope of the moment of inertia from the rotation frequency can change to negative. The latter, by analogy with the terms "backbending" and "upbending", can be called downbending.

Calculations on this basis belong to semi-phenomenological approaches.

In [8], a novel modification is introduced, extending the Arima coefficient to the third order. The computed outcomes of the rotational bands of ²⁴⁴Pu and ²⁴⁸Cm demonstrate an exceptional degree of agreement with experimental observations. Moreover, the used parameterization of the Arima term and its modification successfully describes such behavior of the moment of inertia from frequency as backbending, upbending, and the downturn (down bending).

Commenting on the considered series of works [5-8], it should be kept in mind that the phenomenon of backbending is traditionally associated with the intersection of bands of different nature, namely, the collective one, built on the ground state, and the one that has a two-quasiparticle pair with a sufficiently high spin at its base.

The upbending phenomenon may be due to the fact that the crossing of bands has not yet occurred completely, but the situation is close to it. In addition, the growth of the slope of the moment of inertia from frequency within certain limits is reproduced within the framework of the traditional, but at the same time complete IBM Hamiltonian.

From a microscopic point of view, as the spin of states increases, the average number of quadrupole bosons increases, and this leads to some weakening of pairing. Accordingly, the introduction of the Arima modification of the IBM Hamiltonian was motivated by this decrease in pairing correlation due to the increase in rotation. At the same time, this leads to an increase in the energy of the quasiparticle vacuum or a decrease in the correspon ding depth.

In this case, with an increase in the spin of the collective state, a decrease in the microscopically calculated energy of the d boson ε_d occurs (the corresponding term also arises when using only the $\hat{Q} \cdot \hat{Q}$ interaction by reducing the last operator to the normal order over bosons). Since IBM assumes that the energy of the boson vacuum is constant, a redistribution of the variable part of the energy of the quasiparticle vacuum is made over two boson parameters ε_d and k_1 (see below (8)). This results in the *d*-boson energy practically ceasing to change, and the energy of the boson vacuum remaining unchanged. This procedure is described in [9–11] and clearly demonstrated in [11].

Thus, the weakening of pairing during the unwinding of the nucleus by the values of the parameters of the boson Hamiltonian is significantly leveled out.

There is one more effect left, this is the weakening of the growth of the moment of inertia with frequency at high spins – downturn or downbending. It can be assumed that the IBM specificity, namely, the finiteness of the maximum number of quadrupole bosons, can reproduce this phenomenon [12] without introducing the Arima modifications to the boson Hamiltonian. Introducing the Arima modification is convenient if we remain in the SU(3) approximation of the IBM limit. In this case, this modification approximates a more complex dynamics of collective motions in the nucleus, described in the general case by the SU(6) group.

In [13], the structure of yrast bands in the transuranium nuclei ²⁴²Pu and ²⁴⁴Pu was investigated within the framework of the projected shell model. This approach is completely microscopic.

The description of rotational bands at ultra-high spins within the framework of covariant density functional theory for the $A \ge 242$ actinides was presented in [14].

The rotational bands properties of plutonium isotopes $^{236-246}$ Pu were studied [15] via projected shell model. The results of the calculated energy levels of the yrast band were compared with experimental data and a good agreement has been found. The crossing between two-quasiparticle (2qp) excited bands and the ground state band (g-band) in the high-spin regions has been analyzed in terms of band diagrams. The upbendings are observed in the kinematic moments of inertia curves for $^{236-246}$ Pu isotopes.

The objectives of this work are defined as:

1) obtaining a precise description of the energies of yrast bands in even-even heavy nuclei in the Harris model;

2) comparison of the behavior of $J(\omega^2)$ in the Harris and IBM models both in its phenomenological aspect and in the aspect of its extension by taking into account high-spin modes;

3) development of a criterion for determining the spin at which the crossing of bands occurs without resorting to microscopic calculations;

4) identification of the causes, conditions and values of the spins of collective states when the downturn or downbending effect occurs.

1. PHENOMENOLOGICAL VARIANT OF THE HARRIS MODEL

We will use the Harris model [1] in its phenomenological aspect. If we assume that the following decomposition is valid

$$\sqrt{I(I+1)} = \omega \left(J_0 + \frac{2}{3} J_1 \omega^2 + \frac{3}{5} J_2 \omega^4 + \frac{4}{7} J_3 \omega^6 + \dots \right), \tag{2}$$

then the effective moment of inertia will have the form

$$J = J_0 + J_1 \omega^2 + J_2 \omega^4 + J_3 \omega^6 + \dots$$
(3)

With fixed values of parameters J_0, J_1, J_2, J_3 , the rotation frequencies are found from the equation

$$\frac{2J_0}{\hbar^2} + \frac{2J_1}{\hbar^4} (\hbar\omega)^2 + \frac{2J_2}{\hbar^6} (\hbar\omega)^4 + \frac{2J_3}{\hbar^8} (\hbar\omega)^6 = \frac{2(2I-1)}{\left(\sqrt{I(I+1)} - \sqrt{(I-2)(I-1)}\right)\hbar\omega}.$$
 (4)

For each value of the spin of state *I*, the value $\omega = \omega_I$ is found. The energies of the states of the main band are found as

$$E_{I} = E_{I-2} + \left(\sqrt{I(I+1)} - \sqrt{(I-2)(I-1)}\right)\hbar\omega,$$
(5)

where $E_0 = 0$.

The parameters that determine the moment of inertia (3) are found by minimizing the value

$$\chi^2 = \sum_{I} (E_I^{(\text{exp})} - E_I)^2$$
 (6)

or standard deviation

$$\sigma = \sqrt{\frac{\sum_{I}^{I_{\rm fit}} (E_I^{\rm (exp)} - E_I)^2}{I_{\rm fit}/2}},$$
(7)

where I_{fit} is the maximum spin of the states and the bands by which the parameters J_0, J_1, J_2, J_3 are determined.

The results obtained in the Harris model will be related to the results in the IBM phenomenology. The parameterization of its Hamiltonian is taken as

$$H_{\rm IBM} = \varepsilon_d \,\,\hat{n}_d + k_1 (d^+ \cdot d^+ ss + {\rm H.c.}) + k_2 \left((d^+ d^+)^{(2)} \cdot ds + {\rm H.c.} \right) + \frac{1}{2} \sum_L C_L (d^+ d^+)^{(L)} \cdot (dd)^{(L)}, \quad (8)$$

H.c. means Hermitian conjugation, the dot between the operators corresponds to the scalar product, the quantities ε_d , k_1 , k_2 , C_0 , C_2 , C_4 are the parameters of the model. The total number of bosons or the maximum number of d bosons will be denoted as Ω . For several nuclei, results will be presented within the framework of the extended microscopic version of IBM, where high-spin excitation modes are also considered, without disclosing its content here, but only making a reference to the work [4].

The next section discusses the energies of the yrast band states of even thorium isotopes in comparison with the results of the Harris model phenomenology, and for several nuclei with IBM.

2. ANALYSIS OF MOMENTS OF INERTIA FOR EVEN ISOTOPES OF Th

Figure 1 presents a visual systematics of the energies of the corresponding states. It is evident that the energies of states with spins $I \leq 10^+$ decrease at first to A = 224 rather quickly, and at large nucleon values, the energies decrease quite smoothly, so that $E(2_1^+) = 48.4$ keV for ²³⁶Th. The minimum energy of states with spins $I \geq 12^+$ is achieved in the ²³²Th nucleus.

In Fig. 2, for all even thorium isotopes considered, the effective moments of inertia obtained from the experimental and calculated energies in the variable moment of inertia model, designated as Harr, are compared. For two nuclei, 220,222 Th, the results of calculations in the IBM microscopic version, extended by high-spin excitation modes up to spins 14⁺, are also given. The parameters determining the values of the moments of inertia are given in Table 1, where the standard deviations of the calculated values of energies from the experimental ones are also given relative to the states for which the parameters were selected, the spins of the states to which the parameters were selected, and the maximum spin in the band.

For the lightest of the thorium isotopes under consideration, 220 Th, backbending is observed starting from spin 10⁺, and since the Harris model does not claim to describe it, the parameters were determined from transitions until the state with 8⁺. In this case, in the theory that explicitly considers the elementary modes of two-quasiparticle excitations with high spins [4, 16], the behavior of the moment of inertia associated with the first backward bend is reproduced. This is demonstrated in Fig. 2 associated with this nucleus. Figure 3 shows the composition of wave functions, whose components include collective states constructed only from *d* bosons corresponding to the lowest



Fig. 1. Energies of states of the yrast band in even isotopes of thorium

quadrupole excitations, as well as components including bosons with high multipolarities. According to this figure, for the 8^+ state, the collective component composed only of *d* bosons is greater than 50%. The second backbending, corresponding to the 18^+ spin, should be attributed to the four-quasiparticle excitation.

For the ²²²Th isotope, no manifestation of the bands crossing is observed. However, the average discrepancy between the calculated energies within the Harris scheme and the experimental ones is significantly larger than for heavier thorium isotopes, as follows from Table 1. The reason for this is that although no backbending is observed in this nucleus, a smooth crossing of bands still occurs, as was shown in [4]. This is demonstrated in Fig. 2, corresponding to ²²²Th, where the moments of inertia obtained on the basis of microscopic calculation are also given. Figure 4 shows the composition of the wave functions. According to this figure, for the 12^+ state, ψ^2 (coll) is approximately equal to 50%. Therefore, according to the Harris method, in addition to calculations for all energy differences up to the state with spin 26^+ , calculations were made taking into account the states up to the state with spin 10^+ . In the first case, the difference between the calculated and experimental values up to spin 8^+ is less than 1 keV, and for 10^+ , it is already 5.5 keV. In the second case, this difference for 2^+ , 4^+ is about 2 keV, but for the rest, up to 14⁺, it is less than 1 keV. Together with the previous thorium isotope, this shows that the Harris energy calculation procedure can be successfully applied if the collective component in the wave function is definitely greater than 50%.

For all nuclei, starting with ²²⁴Th, there achieved such an accuracy of reproduction of experimental energies that the difference between the





Nucleus	$\frac{2J_0/\hbar^2}{\mathrm{MeV}^{-1}},$	$2J_1/\hbar^4$, MeV ⁻³	${2J_2/\hbar^6\over MeV^{-5}},$	$2J_3/\hbar^8, MeV^{-7}$	σ , keV	I_{fit}	I_{\max}
²²⁰ Th	-78.027	5627.16	-100806	1032727	0.14E-02	8	22
²²² Th	11.116 17.465	2788.26 3172.00	865.45 -85959.8	0 1995659.8	9.59 0.172	26 10	$\frac{26}{26}$
²²⁴ Th	57.965	2058.09	-7175.0	9656.4	0.33	18	18
²²⁶ Th	82.233	1369.23	4017.6	15137.6	0.33	20	20
²²⁸ Th	102.731	1474.68	-12333.3	349536.8	0.59	22	22
²³⁰ Th	112.141	1038.92	-2054.2	94199.8	0.175	24	24
²³² Th	121.107	928.66	10424.0	-50092.3	0.89	30	30
²³⁴ Th	120.341	984.69	-10471.4	212984.7	0.63	24	24
²³⁶ Th	123.400	844.93	772.3	0	0.18	10	10
²²⁶ U	71.87581	1598.41785	6953.44873	-118350.98438	0.294	14	14
²³⁰ U	115.33965	1119.44153	-3171.23999	121410.60156	0.262	22	22
²³² U	125.72275	944.34747	3252.86279	64539.00781	0.124	20	20
²³⁴ U	137.59946	1030.29688	14549.98926	-126356.42969	0.35	30	30
²³⁶ U	133.45268 132.30104	385.92599 781.21796	19432.23242 -1048.02319	-89088.77344 172107.54688	1.90 0.155	30 24	30 30
²³⁸ U	133.00563	941.80017	-9015.27832	267665.18750	0.468	28	34
²⁴⁰ U	132.39108	388.02209	14673.71387	0.00000	0.413	12	12

Table 1. Parameters determining the effective moments of inertia for Th and U isotopes





Fig. 3. The composition of the wave functions of the states of the yrast band obtained on the basis of the microscopic model for $^{220}\mathrm{Th}$

Fig. 4. The composition of the wave functions of the states of the yrast band obtained on the basis of the microscopic model for $^{222}\mathrm{Th}$

calculated energies and the experimental ones does not exceed 1 keV on average, as can be seen from the σ values given in the same Table 1. This corresponds to the work [4], where it was found that in thorium isotopes, starting with this one, in all observed states of yrast bands, the collective

component remains the main component. Table 2 presents the experimental energies of states and the difference between the calculated and experimental energies for $^{220-224}$ Th. For $^{226,\,228}$ Th nuclei, a comparison of the experimental and calculated ones is given in Table 3, for $^{230,\,232}$ Th — in Table 4, for $^{234,\,236}$ Th — in Table 5. In almost all of these nuclei, the difference between the

Table 2.	Comparison of	experimental	[17] and	theoretical	energy	values	in l	keV
	-	for ^{220,222,2}	²²⁴ Th nu	clei				

I^{π}	220	Th	22	²² Th	22	⁴ Th
1	Exp.	$E_{\rm cal} - E_{\rm exp}$	Exp.	$E_{\rm cal} - E_{\rm exp}$	Exp.	$E_{\rm cal} - E_{\rm exp}$
2^{+}	386.5(1)	-0.001	182.9(2)	-0.06	98.1(3)	-0.12
4^{+}	759.80(15)	0.002	439.2(3)	0.22	284.1(5)	0.58
6^{+}	1166.03(17)	-0.002	749.3(4)	-0.29	534.7(5)	-0.30
8+	1598.16(20)	-0.0005	1092.8(5)	0.09	833.9(6)	-0.46
10^{+}	2012.73(23)	38	1460.8(5)	-0.01	1173.8(6)	-0.01
12^{+}	2441.9(3)	78	1850.6(5)	-3.8	1549.8(6)	0.43
14^{+}	2885.0(3)	118	2259.7(6)	-12	1958.9(7)	0.21
16^{+}	3376.4(6)	121	2688.0(6)	-28	2398.0(7)	-0.27
18^{+}	3867.1(6)	136	3133.9(6)	-51	2864	0.01
20^{+}	4319.6(7)	198	3596.8(7)	-82		
22^{+}	4716.1(12)	324	4078.6(7)	-124		
24^{+}			4579.2(7)	-177		
26^{+}			5099.2(9)	-244		

Table 3. Comparison of experimental [17] and theoretical energy values in keV for 226,228 Th nuclei

I^{π}	22	⁶ Th	228	Th
1	Exp.	$E_{\rm cal} - E_{\rm exp}$	Exp.	$E_{\rm cal} - E_{\rm exp}$
2^{+}	72.20(4)	-0.27	57.773(3)	0.17
4^{+}	226.43(5)	0.36	186.838(3)	0.08
6^+	447.3(2)	0.69	378.195(12)	-0.46
8+	721.9(2)	0.38	622.5(3)	-0.85
10^{+}	1040.3(3)	-0.10	911.8(3)	-0.78
12^{+}	1395.2(4)	-0.24	1239.3(4)	-0.21
14^{+}	1781.5(5)	0.02	1599.4(5)	0.55
16^{+}	2195.8(6)	0.22	1987.9(6)	0.64
18^{+}	2635.1(7)	0.32	2400.5(8)	0.23
20^{+}	3097.1(8)	0.15	2834.4	-1.2
22^{+}			3283.4	-0.15

I^{π}	230	Th	232	²Th
1	Exp.	$E_{\rm cal} - E_{\rm exp}$	Exp.	$E_{\rm cal} - E_{\rm exp}$
2^{+}	53.230(11)	0.04	49.369(9)	0.02
4^{+}	174.119(17)	0.019	162.12(2)	0.08
6^{+}	356.54(12)	0.01	333.26(8)	0.29
8+	593.89(17)	-0.27	556.9(1)	0.23
10^{+}	879.36(24)	-0.39	826.8(1)	-0.22
12^{+}	1206.7(5)	0.06	1137.1(5)	-0.95
14^{+}	1571.9(6)	-0.05	1482.2(6)	-1.1
16^{+}	1969.6(7)	0.13	1858.5(7)	-1.2
18^{+}	2396.4(9)	0.09	2262.4(9)	-0.56
20^{+}	2848.7(10)	0.09	2691(1)	0.95
22^{+}	3324.1(11)	-0.32	3144(1)	1.5
24^{+}	3819.1(15)	-0.05	3620.0(15)	0.89
26^{+}			4117(2)	-0.50
28^{+}			4633(2)	-1.8
30^{+}			5164(3)	-0.15

Table 4. Comparison of experimental [17] and theoretical energy values in keV for 230,232 Th nuclei

Table 5. Comparison of experimental [17] and theoretical energy values in keV for 234,236 Th nuclei

I^{π}	234	¹ Th	2	³⁶ Th
1	Exp.	$E_{\rm cal} - E_{\rm exp}$	Exp.	$E_{\rm cal} - E_{\rm exp}$
2^{+}	49.55(6)	0.14	48.4(3)	0.09
4^{+}	163.05(12)	0.14	160.0(6)	-0.36
6^{+}	336.45(24)	-0.24	329.4(7)	0.14
8+	564.7(3)	-0.80	553.4(7)	-0.06
10^{+}	842.5(4)	-1.03	826.1(9)	-0.04
12^{+}	1164.9(6)	-0.81		
14^{+}	1526.6(7)	0.11		
16^{+}	1923.4(8)	0.80		
18^{+}	2351.0(9)	0.72		
20^{+}	2805.1(11)	-0.13		
22^{+}	3281.4(12)	-1.04		
24^{+}	3775.1(13)	-0.15		

calculated energies and the experimental ones does not exceed 1 keV, and for the $^{224,\,236}\text{Th}$ nuclei, the calculated energy values differ from the experimental ones by a smaller amount than the experimental errors. Therefore, anomalies in the behavior of $J(\omega^2)$ for ^{236}Th in Fig.2 should be attributed to the

experimental errors. This anomaly will disappear if the experimental value $E(4^+) = 160$ keV is reduced by a value from 0.358 to 0.37 keV. Such a high quality of description of the energies of states in the model of variable moment of inertia in heavy nuclei is achieved given that excitation energies can reach up to 5 MeV, and the spin of states — up to 30^+ .

Heavy nuclei in the majority have a stable deformation, and the deviation to the smaller side of the energies in the band from the dependence I(I + 1)has a number of reasons, partly discussed earlier. This corresponds to the growth of the moment of inertia from the rotation frequency. The monotony of such a dependence at sufficiently large spins of states can be violated, leading, for example, to backbending, and this invariably occurs in medium nuclei. Also, an anomaly in the behavior of the moment of inertia can manifest itself through upbending, when the intersection of bands is not fully completed. And finally, downbending. This effect can manifest itself at spins of at least 26^+ and in those cases when, for example, a state with the same spin, but in the S band, turns out to be significantly higher and the observed state remains mainly collective. Moreover, the energies of such high-spin collective states may turn out to be somewhat higher than those that would be obtained based on the trends corresponding to lower spins. There may be two reasons for such a behavior of the energies of collective states at high spins. One is associated with the reduction of the configuration space of two-quasiparticle states that form the D phonon, in the presence of multibosons, which is realized to a greater extent at high spins. To quantitatively evaluate this phenomenon at the energy of collective states, it is necessary to carry out self-consistent solutions for each of the collective states separately, as was done for medium nuclei [9]. However, for heavy nuclei, such a procedure is difficult to implement due to the high sensitivity of the calculated energies to the IBM Hamiltonian parameters. Another reason for some additional growth of energies of collective states at high spins, which is what downbending gives, may be the specificity of the description of collective states in the general case of the SU(6) symmetry, which consists in the finiteness of the maximum number of quadrupole bosons Ω .

In the variable moment of inertia model, the growth of the slope $J(\omega^2)$ is effectively reproduced. One of the reasons for this growth is the gradual increase of the noncollective component in the states of the yrast band. This explains why in deformed nuclei the quality of the description of energies in the Harris model is better than in the IBM phenomenology. The situation regarding IBM changes if we move from its phenomenology to the microscopic version, when the IBM parameters are calculated and high-spin excitation modes are explicitly taken into account. This also allows us to determine the limits of applicability of the Harris scheme in the region of spins where the bands cross. For example, a microscopic calculation of the structure of the states of the yrast band in ²²²Th showed [4] that the Harris scheme successfully reproduces collective states in which the collective component is noticeably larger than any of the others, and itself can be somewhat less than 50%.

3. ANALYSIS OF MOMENTS OF INERTIA FOR EVEN ISOTOPES OF U

The parameters determining the moment of inertia for U isotopes are given in Table 1. The corresponding moments of inertia are given in Fig. 5. In addition to the experimental and calculated values of the moments of inertia according to Harris, the results of calculations in accordance with the IBM phenomenology are given, excluding high-spin modes.

The parameters of the boson Hamiltonian (8) for the U isotopes are given in Table 6 and the corresponding moments of inertia with others are also given in Fig. 5. The parameters of the boson Hamiltonian given in Table 6 are given with an accuracy of 1 eV. To justify the need for such high accuracy, the partial derivatives $\partial E_I / \partial \varepsilon_d, \ldots, \partial E_I / \partial C_4$ were calculated in the vicinity of the parameter values determined for ²³⁶U. They are given in Table 7. In accordance with these values, the maximum deviations of the parameters were obtained when the deviations of the calculated energies do not exceed 0.1 keV. These values are given in Table 8. The change in the Hamiltonian parameter (8) k_1 reacts most strongly to the energies of the states under consideration. And as can be seen from Table 8, at high spins its values should be given with greater accuracy than 1 eV, but since at high spins the experimental uncertainty of energies exceeds 1 keV, this can be avoided for now.

The IBM Hamiltonian parameters, including the Ω number, for $^{226-238}$ U were determined from states up to the limiting observed spins and are given in Table 6. Let us discuss the quality of the description of the energies of states consistently in all uranium isotopes under consideration.

Nucleus	ε_d	k_1	k_2	C_0	C_2	C_4	Ω
²²⁶ U	-0.080461	-0.030253	0.009227	0.005134	0.060668	0.001103	20
²²⁶ U	0.036599	-0.038003	0.004841	-0.010224	0.033211	-0.034492	28
²²⁶ U	0.040505	-0.036817	0.004122	0.136447	0.035798	-0.031713	34
²²⁶ U	-0.059896	-0.037406	0.004168	0.001010	0.011875	-0.020448	25
²³⁰ U	-0.614828	-0.058595	0.032967	0.899961	0.061097	0.045202	21
²³² U	-0.725962	-0.058227	0.032505	0.739341	0.034253	0.054798	19
²³⁴ U	-0.594347	-0.049232	0.032013	0.674469	0.076536	0.039112	22
²³⁴ U	-0.580119	-0.049519	0.034100	0.637994	0.071404	0.038430	21
²³⁶ U	-0.626829	-0.059598	0.051147	0.713640	0.042979	0.019237	25
²³⁸ U	-0.634858	-0.060349	0.053701	0.687343	0.039218	0.016058	25
²⁴⁰ U	-0.634453	-0.059500	0.050521	0.717118	0.029825	0.022396	25

Table 6. IBM Hamiltonian parameters for even U nuclei





for 236 U, E_2 and below are understood to be the excitation energy, and E_0 – the total boson energy, measured from the Table 7. Partial derivatives of IBM energies with respect to boson parameters in the special case of parameters adopted energy of the *d*-boson vacuum

+ 30+	9 7.7	16 - 234	13 -9.5	.4 -8.2	7 7.3	.2 104
+ 28	.2 6.	96 -2	5.2 -1	6.6 -7	.0 6.	.7 90
+ 26	4 6.	75 -1	5.3 -1	<u>8</u> .	3	2 77
- 24		-1-	.6 -16	1 -5	20	5 66.
- 22-	4.7	4 -15	3 -16	2 -5.	4.6	0 55.
- 20+	4.0	3 -13	-10	6 -4.	3.9	4 46.0
- 184	3.3	.7 -11	.4 -1.	9 -3.	3.3	5 37.4
16^{+}	2.7	8 -93.	4 -13	-2.	2.7	29.6
14^{+}	2.2	1 -74.	-11.	-2.5	2.2	22.9
12^{+}	1.7	-57.4	-9.3	-1.7	1.7	17.0
10^{+}	1.2	-41.7	-7.1	-1.21	1.19	11.9
8+	0.8	-28	-5.0	-0.8	0.8	7.8
$^{+9}$	0.48	-16.7	-3.09	-0.47	0.48	4.55
$^{++}$	0.23	-8.1	-1.54	-0.23	0.23	2.16
2^{+}	0.071	-2.5	0.475	-0.067	0.071	0.65
+0	11	316	-138	12	-17	23
I^{π}	$\partial E_I/\partial \varepsilon_d$	$\partial E_I / \partial k_1$	$\partial E_I / \partial k_2$	$\partial E_I / \partial C_0$	$\partial E_I / \partial C_2$	$\partial E_I / \partial C_4$

Table 8. The required accuracy of the parameters in eV so that the accuracy of calculating the energy of the yrast band states is not less than 0.1 keV in the particular case of the parameters adopted for 236 U, for $I \neq 0$, the energies are considered relative to the ground state

I^{π}	$^{+0}$	2^{+}	$^{++}$	$^{+9}$	*	10^{+}	12^{+}	14^{+}	16^{+}	18^{+}	20^{+}	22^{+}	24^{+}	26^{+}	28^{+}	30^{+}
$\Delta \varepsilon_d$	6	1400	430	210	125	83	59	45	37	30	25	21	18	10	14	13
$ riangle k_1$	0.32	40	12	9	3.6	2.4	1.7	1.3	1.1	0.88	0.75	0.65	0.57	0.51	0.46	0.43
$ riangle k_2$	0.72	210	65	32	20	14	11	8.8	7.5	6.7	6.2	6.0	6.1	6.6	7.7	10
$\triangle C_0$	8	1492	435	213	125	83	59	45	34	28	24	20	17	15	14	12
$\triangle C_2$	9	1408	435	208	125	84	59	45	37	30	26	22	19	17	15	14
$\triangle C_4$	4	154	46	22	13	8.4	5.9	4.4	3.4	2.7	2.2	1.8	1.5	1.3	1.5	1.3

For ²²⁶U, the experimental data show a practically linear dependence of $J(\omega^2)$ and this is naturally reproduced in the Harris scheme. This could not be obtained in the description in IBM. The calculations were carried out in a wide range of Ω from 14 to 34. In this case, if the quality of reproduction of energies for states with $I \leq 12^+$ is satisfactory, and this is evident from Table 9, where the values of energies with $\Omega = 20$ are given, then the energy of the state with 14^+ turns out to be underestimated by 7 keV, and this leads to the calculated increase in the value of J. If we degrade the quality of the energy description, the linear dependence $J(\omega^2)$ is approximately reproduced, which is reflected in the corresponding Fig.5. The question of the possibility of quantitative reproduction of the energies of the ²²⁶U nucleus in IBM remains open. This is despite the fact that the question concerns the description of states up to relatively low spins. It should be borne in mind that the experimental errors for this isotope are the highest of all those considered.

Table 9. Comparison of experimental [17] and theoretical energy values in keV for 226 U nuclei; for IBM, $\Omega=20$

I^{π}			²²⁶ U		
1	Exp.	$E_{\rm cal}$	$E_{\rm cal} - E_{\rm exp}$	$E_{\rm IBM}$	$E_{\rm IBM} - E_{\rm exp}$
2^{+}	81.3(6)	81.464	0.16	81.510	0.21
4^{+}	250.0(9)	249.682	-0.32	248.46	-1.54
6^+	483.2(9)	482.79	-0.41	482.24	-0.92
8+	766.4(10)	766.51	0.11	768.45	2.05
10^{+}	1091.6(10)	1091.98	0.38	1095.9	4.3
12^{+}	1453.8(10)	1453.42	-0.38	1455.7	1.9
14^{+}	1847.0(13)	1846.93	-0.07	1840.1	-6.9

For the next nucleus, 230 U, the description of energies in the Harris scheme is precise and the difference between the calculated and experimental values of energies, as can be seen from Table 10, does not exceed 0.4 keV. Within the IBM framework, this is 1.4 keV, which also indicates a high quality of the description. This is despite the fact that states are observed up to 22^+ .

For 232 U, states are observed up to 20^+ and the description of their energies in the two calculations is quite satisfactory. Thus, in the Harris scheme, the discrepancy is no more than 0.21 keV, and for IBM — no more than 1.2 keV, as follows from Table 11. The quality of the energy description is especially clearly manifested through the moments of inertia for 230 U and 232 U in Fig. 5.

For ²³⁴U, states are observed already up to 30^+ . In the Harris scheme, the discrepancy is no more than 0.86 keV. Within the IBM framework, as can be seen from Fig. 5, calculations were made with two values of Ω , these are 21 and 22. The results of the calculations of the moments of inertia closely cover the moments of inertia from the experimental energies on both sides. Taking into account the influence of high-spin modes on the energies of states,

I^{π}			²³⁰ U		
1	Exp.	$E_{\rm cal}$	$E_{\rm cal} - E_{\rm exp}$	$E_{\rm IBM}$	$E_{\rm IBM} - E_{\rm exp}$
2^{+}	51.737(23)	51.797	0.060	51.623	-0.114
4^{+}	169.35(4)	169.356	0.006	169.17	-0.18
6^+	346.96(20)	346.85	-0.11	347.04	0.08
8+	578.1(3)	577.76	-0.34	578.59	0.49
10^{+}	856.3(3)	855.91	-0.39	857.24	0.94
12^{+}	1175.6(4)	1175.68	0.08	1177.0	1.40
14^{+}	1531.5(4)	1532.07	0.57	1532.7	1.20
16^{+}	1921.0(5)	1920.65	-0.35	1920.1	-0.90
18^{+}	2337.6(6)	2337.56	-0.04	2335.9	-1.7
20^{+}	2779.6(11)	2779.47	-0.13	2777.6	-2.0
22^{+}	3243.6(15)	3243.54	-0.06	3243.7	0.1

Table 10. Comparison of experimental [17] and theoretical energy values in keV for 230 U nuclei; for IBM, $\Omega = 21$

Table 11. Comparison of experimental [17] and theoretical energy values in keV for 232 U nuclei; for IBM, $\Omega = 19$

I^{π}			²³² U		
1	Exp.	$E_{\rm cal}$	$E_{\rm cal} - E_{\rm exp}$	$E_{\rm IBM}$	$E_{\rm IBM} - E_{\rm exp}$
2^{+}	47.573(8)	47.590	0.017	47.533	-0.04
4^{+}	156.566(10)	156.545	-0.021	156.49	-0.076
6^+	322.69(7)	322.76	0.065	322.80	0.11
8+	541.1(1)	540.97	-0.13	541.23	0.13
10^{+}	805.88(16)	805.68	-0.20	806.16	0.28
12^{+}	1111.6(2)	1111.59	-0.01	1112.1	0.50
14^{+}	1453.8(3)	1453.91	0.10	1454.1	0.30
16^{+}	1828.2(4)	1828.39	0.19	1828.0	-0.20
18^{+}	2231.6(6)	2231.39	-0.21	2230.4	-1.2
20^{+}	2659.8(9)	2659.78	-0.02	2659.3	-0.5

we give preference to $\Omega = 21$ and it is with this value that the calculated values are presented in Table 12, from which it is clear that the corresponding differences are no more than 3.2 keV.

In the ²³⁶U nucleus, the observed states also extend to $I = 30^+$. The observed moments of inertia at $I = 28^+$ and $I = 30^+$ definitely give a decrease in the slope of $J(\omega^2)$. More precisely, starting from spin 26⁺, a slower growth of the moment of inertia — downbending — is manifested. That is why the parameters in the Harris scheme were determined by the states up to spin $I = 24^+$ and for these states, the difference between the experimental and

I^{π}	²³⁴ U				
1	Exp.	$E_{\rm cal}$	$E_{\rm cal} - E_{\rm exp}$	$E_{\rm IBM}$	$E_{\rm IBM} - E_{\rm exp}$
2^{+}	43.4981(10)	43.501	0.003	43.517	-0.195
4^{+}	143.352(4)	143.362	0.010	143.34	1.92
6^{+}	296.072(4)	296.12	0.046	295.89	0.97
8+	497.04(3)	497.06	0.02	496.63	0.17
10^{+}	741.2(5)	741.13	-0.07	740.75	0.54
12^{+}	1023.8(7)	1023.56	-0.24	1023.6	0.60
14^{+}	1340.5(12)	1340.22	-0.28	1340.9	1.50
16^{+}	1687.8(16)	1687.70	-0.1	1689.1	1.90
18^{+}	2062.8(17)	2063.20	0.4	2064.9	2.20
20^{+}	2464.0(18)	2464.49	0.49	2465.9	2.30
22^{+}	2889.5(18)	2889.78	0.28	2890.3	2.0
24^{+}	3338.5(21)	3337.64	-0.86	3336.9	1.9
26^{+}	3807.5(23)	3806.98	-0.52	3805.0	3.2
28^{+}	4296.5(25)	4296.95	0.45	4294.8	3.2
30+	4807	4806.98	-0.02	4807.1	3.2

Table 12. Comparison of experimental [17] and theoretical energy values in keV for 234 U nuclei; for IBM, $\Omega = 21$

Table 13. Comparison of experimental [17] and theoretical energy values in keV for 236 U nuclei; for IBM, $\Omega=25$

I^{π}	²³⁶ U				
1	Exp.	$E_{\rm cal}$	$E_{\rm cal} - E_{\rm exp}$	$E_{\rm IBM}$	$E_{\rm IBM} - E_{\rm exp}$
2^{+}	45.2431(20)	44.915	-0.33	45.204	-0.0391
4^{+}	149.480(5)	148.921	-0.56	149.43	-0.05
6^+	309.788(6)	309.85	0.062	309.87	0.082
8^{+}	522.26(4)	523.77	1.5	522.58	0.32
10^{+}	782.4(5)	785.39	2.99	782.93	0.53
12^{+}	1085.4(7)	1088.97	3.6	1086.0	0.6
14^{+}	1426.4(9)	1429.25	2.8	1426.8	0.4
16^{+}	1801.0(10)	1801.74	0.74	1800.8	-0.2
18^{+}	2204.0(12)	2202.78	-1.2	2203.7	-0.3
20^{+}	2631.8(13)	2629.43	-2.4	2631.7	-0.1
22^{+}	3081.0(14)	3079.34	-1.7	3081.9	0.9
24^{+}	3550.0(17)	3550.57	0.57	3552.0	2.0
26^{+}	4039.0(20)	4041.55	2.6	4040.9	1.9
28^{+}	4549.0(22)	4550.99	1.99	4548.4	-0.6
30^{+}	5077(4)	5077.80	0.8	5075.8	-1.2

calculated energies does not exceed 0.74 keV, as follows from Table 13. The moments of inertia at high spins, as can be seen from Fig.5, for this nucleus significantly exceed the experimental values. The calculation in IBM

reproduces this experimental trend and this is achieved at $\Omega = 25$. This can be associated with a decrease in the collective configuration space at the corresponding spins, as well as the presence of *s* bosons or square roots, due to which the closure of the *SU*(6) algebra of boson operators is realized in comparison with other collective models. This is the case when the variable moment of inertia model with the used parameterization cannot describe this phenomenon, and the IBM phenomenology describes the corresponding increase in the energies of collective states. Such a phenomenon is realized, as can be judged by ^{236, 238}U nuclei, starting with spin 28⁺. For all states of this nucleus, the difference between the calculated IBM and experimental energy values does not exceed 2 keV.

The effect observed in the ²³⁶U nucleus of weakening the growth of J from ω^2 at $I = 28^+$, 30^+ is even more pronounced in the ²³⁸U nucleus, since the spins observed for it extend to $I = 34^+$. The parameters in the Harris scheme were determined by the states with $I \leq 28^+$ and for these states, as can be seen from Table 14, the difference between the experimental and calculated energies does not exceed 0.76 keV. For large spins, this difference grows rapidly, and the calculated energy values are noticeably lower than the experimental ones. As can be seen from Fig. 5, for this nucleus, the calculation in IBM reproduces the experimental situation and this is realized at $\Omega = 25$. If we use large values of Ω , for example, 30, then J will be significantly larger, approaching those given by the Harris scheme. The difference between the

I^{π}	²³⁸ U				
1	Exp.	$E_{\rm cal}$	$E_{\rm cal} - E_{\rm exp}$	$E_{\rm IBM}$	$E_{\rm IBM} - E_{\rm exp}$
2^{+}	44.916(13)	45.004	0.0883	44.785	-0.131
4^{+}	148.38(3)	148.394	0.014	148.10	-0.28
6^{+}	307.18(8)	307.12	-0.057	307.26	0.08
8+	518.1(3)	517.45	-0.65	518.49	0.31
10^{+}	775.9(4)	775.32	-0.58	777.25	1.35
12^{+}	1076.7(5)	1076.38	-0.32	1078.7	2.0
14^{+}	1415.5(6)	1415.94	0.44	1417.9	2.4
16^{+}	1788.4(6)	1789.15	0.75	1790.1	1.7
18^{+}	2191.1(7)	2191.41	0.31	2191.0	-0.1
20^{+}	2619.1(8)	2618.62	-0.48	2616.7	-2.4
22^{+}	3068.1(9)	3067.34	-0.76	3064.1	-4.1
24^{+}	3535.3(12)	3534.76	-0.54	3530.7	-4.6
26^{+}	4018.1(16)	4018.57	0.47	4015.1	-3.0
28^{+}	4517	4516.92	-0.084	4517.1	0.1
30^{+}	5035.1(21)	5028.27	-6.83	5038.0	2.9
32^{+}	5581(3)	5551.37	-29.6	5580.7	-0.3
34^{+}	6146(4)	6085.16	-60.8	6149.9	3.9

Table 14. Comparison of experimental [17] and theoretical energy values in keV for 238 U nuclei; for IBM, $\Omega = 25$

experimental and calculated energies within the IBM framework for all spins does not exceed 4.6 keV.

It should be noted that the description of the energies of states in a number of uranium nuclei is so good that the discrepancies with experimental values sometimes give smaller values than the experimental uncertainties, especially for calculations in the Harris scheme.

As a rule, the quality of description in the Harris model is higher than in IBM, this is evident from ^{226,230,232,234}U (there are no experimental data for ²²⁸U). This can be explained by the method of determining the parameters. For the Harris scheme, they are determined by the least squares method, which makes them clearly optimal, while for IBM, other methods for determining the parameters have to be used, which are not so effective. For ²³⁶U, as can be seen from Fig. 5, both models reproduce the energies equally qualitatively.

For ²⁴⁰U, states are known only up to spin 12⁺. In Fig.5, for ²⁴⁰U, a strong anomaly of the moment of inertia for the first excitation is visible. It cannot be reproduced by any of the models and its presence is associated with a large experimental error in the energy of the first excitation (see Table 15). If, when describing energies within the IBM framework, the determination of parameters is carried out without the 2^+_1 state, then its energy turns out to be equal to 45.6 keV, that is, 0.6 keV higher than that suggested by the experiment. In this case, the experimental error for this state is equal to 1 keV, i.e., such a correction is within the experiment of inertia is the result of a large experimental uncertainty. If the moments of inertia are determined with the corrected energy of the 2^+ state, the value of σ will decrease significantly and become equal to 0.08.

I^{π}	²⁴⁰ U				
	Exp.	$E_{\rm cal}$	$E_{\rm cal} - E_{\rm exp}$	$E_{\rm IBM}$	$E_{\rm IBM} - E_{\rm exp}$
2^{+}	45(1)	45.274	0.27	45.599	0.6
4^{+}	150.60(10)	150.112	-0.49	150.80	0.2
6^{+}	313.19(14)	312.41	-0.78	312.91	-0.28
8+	528.69(18)	528.43	-0.26	528.12	-0.57
10^{+}	792.9(3)	793.00	0.1	791.95	-0.95
12^{+}	1100.5(4)	1100.32	-0.18	1099.6	-0.9

Table 15. Comparison of experimental [17] and theoretical energy values in keV for 240 U nuclei; for IBM, $\Omega=25$

CONCLUSIONS

A comparison of the effective moments of inertia for heavy even nuclei, namely, thorium and uranium isotopes, with the variable moment of inertia model and IBM, both in its phenomenological aspect and in its microscopic version, allowed us to draw the following conclusions.

1. The Harris model effectively reproduces the experimental situation if the collective component in the wave function remains at least 50%. The standard boson model cannot do this; for this, it must be extended by explicitly taking into account high-spin excitation modes up to $J \leq 14^+$. However, the Harris scheme does not reproduce the decrease in the slope of J from ω^2 .

2. If the energies of collective states increase significantly at spins greater than 28^+ , this may be due to a reduction in the collective configuration space. This situation is reproduced in the IBM phenomenology, but cannot be reproduced in either the Harris model or classical geometric collective models.

In this paper, the Harris scheme is not considered through the use of the cracking model, but only as one of the convenient and visual representations of the excitation energies in the bands. In a number of cases for the lowest states, when there are large experimental uncertainties, it can give clarifying estimates. For large values of spins, when the experimental values of the moments of inertia are noticeably greater than the Harris estimates, it is possible to make judgments about the extent of the influence of noncollective modes with high spins on these states.

REFERENCES

- 1. Harris S. M. // Phys. Rev. B. 1965. V. 138. P. 509.
- 2. Efimov A. D., Izosimov I. N. // Phys. At. Nucl. 2021. V. 84. P. 660.
- 3. Arima A., Iachello F. // Phys. Rev. Lett. 1975. V. 35. P. 1069.
- 4. Efimov A. D., Izosimov I. N. JINR Preprint E4-2024-27. Dubna, 2024.
- 5. Yoshida N., Sagawa H., Otsuka T., Arima A. // Phys. Lett. B. 1991. V.256. P. 129.
- 6. Liu Y.-X., Song J.-G., Sun H.-Z., Zhao E.-G. // Phys. Rev. C. 1997. V. 56. P. 1370.
- 7. Liu Y., Song J., Sun H. Zhou, Wang J. Jun, Zhao E.-G. // J. Phys. G: Nucl. Part. Phys. 1998. V.24. P. 117.
- 8. Dadwal A., He X.-T. arXiv: 2403.05823v1 [nucl-th]. 2024.
- 9. Efimov A. D. // Phys. At. Nucl. 2020. V. 83. P. 380.
- 10. Efimov A. D., Mikhajlov V. M. // Bull. Russ. Acad. Sci. 2012. V. 76. P. 857.
- 11. Efimov A. D., Mikhajlov V. M. // Bull. Russ. Acad. Sci. 2011. V. 75. P. 890.
- 12. Efimov A.D., Koval I.V., Izosimov I.N., Usmanov P.N. JINR Preprint E4-2024-61. Dubna, 2024.
- 13. Yue Z., Zhong K. Xu., Shui-Fa S., Yu-Peng Y., Chuang-Ye H., Shi-Wei Y. // Chin. Phys. Lett. 2012. V. 29. P. 052101.
- 14. Afanasjev A. V. // Phys. Scr. 2014. V. 89. P. 054001.
- 15. Jaafer F. A., Al-Khudair F. H. // Phys. Rev. C. 2022. V. 106. P. 064309.
- 16. Efimov A. D., Izosimov I. N. // Phys. At. Nucl. 2023. V. 86. P. 333.
- 17. National Nuclear Data Center, Brookhaven National Laboratory. http://www.nndc.bnl.gov.

Received on February 5, 2025.

Редактор В. В. Булатова

Подписано в печать 05.03.2025. Формат 60 × 90/16. Бумага офсетная. Печать цифровая. Усл. печ. л. 1,5. Уч.-изд. л. 1,71. Тираж 100 экз. Заказ № 61056.

Издательский отдел Объединенного института ядерных исследований 141980, г. Дубна, Московская обл., ул. Жолио-Кюри, 6. E-mail: publish@jinr.ru www.jinr.ru/publish/